Author Index

Aburatani, R., see Okamoto, Y. 454
Adachi, S., see Hashimoto, K. 241
Al-Khalidi, A. H. E., see Girard, J. E. 355
Altmayer, B., see Schwenk, S. 424
Amato, A., see Gagliardi, L. 296
Andreolini, F., see Flurer, C. L. 73
Arai, T.
——, Koike, H., Hirata, K. and Ōizumi, H.
Separation of pyridone carboxylic acid enantiomers by high-performance liquid chromatography using copper(II)–L-amino acid as the eluent 439
Araki, M., see Tanaka, N. 95
Armstrong, D. W.
——, He, F.-Y. and Han, S. M.
Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 345
Beasly, V. R., see Harada, K.-I. 275
Beger, J., see Voelkel, A. 219
Benická, E., see Krupčik, J. 203
Bezard, J., see Narce, M. 249
Biedermann, M., see Grob, K. 387
Borra, C., see Flurer, C. L. 73
Buell, P. E., see Girard, J. E. 355
Caponecchi, G., see Polcaro, C. 127
Carmichael, W. W., see Harada, K.-I. 275
Caude, M., see Lienne, M. 55
Cavazzutti, G., see Gagliardi, L. 296
Chiou, S.-H.
——, and Wang, K.-T.
Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 404
Cho, S. Y., see Nam, H. S. 445
Cohen, A. S., see Guttman, A. 41
Cozzi, R., see Polcaro, C. 127
Crouch, Jr., F. W.
——, Riley, C. M. and Stobaugh, J. F.
High-performance liquid chromatography of bis[1,2-bis-(diphenylphosphino)ethane]gold(I) chloride, a potential antineoplastic agent 333
Dahiya, J. S.
—— and Rimmer, S. R.
High-performance liquid chromatography of phytoalexins in stem callus tissues of rapeseed 448
Dahlem, A. M., see Harada, K.-I. 275
Eichhorn, K. W., see Schwenk, S. 424
Fiore, M., see Polcaro, C. 127
Flurer, C. L.
——, Borra, C., Andreolini, F. and Novotny, M.
Separation of proteins by microcolumn liquid chromatography based on the reversed-phase and size-exclusion principles 73
Fujii, Y.
——, Ikeda, Y. and Yamazaki, M.
Micro high-performance liquid chromatographic determination of cardiac glycosides in β-methyldigoxin and digoxin tablets 157
Gagliardi, L.
——, Amato, A., Cavazzutti, G., Tonelli, D. and Montanarella, L.
Identification of xanthene dyes in lipsticks by reversed-phase high-performance liquid chromatography 296
Gál, S., see Kiss, L. 109
Galensa, R., see Risch, B. 291
Gangoda, M. E.
—— and Gilpin, R. K.
Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase liquid chromatographic modifiers 365
Gawdzik, B.
—— and Patrykiejew, A.
Gas chromatography on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)-naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 233
Gilpin, R. K., see Gangoda, M. E. 365
Girard, J. E.
——, Rebbani, N., Buell, P. E. and Al-Khalidi, A. H. E.
Carbohydrate–borate eluents for anion chromatography 355
Gorbunov, A. A.
——, Solovyova, L. Ya. and Pasechnik, V. A.
Fundamentals of the theory and practice of polymer gel-permeation chromatography as a method of chromatographic porosimetry 307
Greenaway, W.
——, Wollenweber, E., Scaysbrook, T. and Whatley, F. R.
Novel isoferulate esters identified by gas chromatography–mass spectrometry in bud exudate of *Populus nigra* 284
Gref, R.
Gas chromatographic analysis of underivatized resin acids 428

Gresti, J., see Narce, M. 249
Grinberg, N., see Guttman, A. 41
Grob, K.
——, Biedermann, M. and Li, Z.
Checking the capacity of a splitless injector — a simple test 387
Günther, K.
Thin-layer chromatographic enantiomeric resolution via ligand exchange 11
Guttman, A.
——, Paulus, A., Cohen, A. S., Grinberg, N. and Karger, B. L.
Use of complexing agents for selective separation in high-performance capillary electrophoresis. Chiral resolution via cyclodextrins incorporated within polyacrylamide gel columns 41
Hagman, A.
—— and Jacobsson, S.
Trapping efficiency of capillary cold traps 117
Han, S. M., see Armstrong, D. W. 345
Harada, K.-I.
——, Matsuura, K., Suzuki, M., Oka, H., Watanabe, M. F., Oishi, S., Dahlem, A. M., Beasly, V. R. and Carmichael, W. W.
Analysis and purification of toxic peptides from cyanobacteria by reversed-phase high-performance liquid chromatography 275
Hashidzume, K., see Tanaka, N. 95
Hashimoto, H.
——, Koyama, Y. and Shimura, T.
Isolation of *cis–trans* isomers of canthaxanthin by high-performance liquid chromatography using a calcium hydroxide column and identification of their configurations by ^{1}H NMR spectroscopy 182
Hashimoto, K.
——, Shirai, Y., Adachi, S. and Horie, M.
Effect of the content of divinylbenzene in ion-exchange resins on the chromatographic separation of α-cyclodextrin and glucose 241
Hatada, K., see Okamoto, Y. 454
He, F.-Y., see Armstrong, D. W. 345
Heimler, D.
——, Vidrich, V.
Thin-layer chromatographic characterization of essential oils 301
Hernádi, F. J., see Kiss, L. 109
Herrmann, K., see Risch, B. 291
Hevesi, T., see Krupčik, J. 203
Hirata, K., see Arai, T. 439
Hoque, E.
Isocratic reversed-phase high-performance liquid chromatographic analysis of pigments in Norway spruce 417
Horie, M., see Hashimoto, K. 241
Hostettmann, K., see Slacanin, I. 265
Ikeda, Y., see Fujii, Y. 157
Iwaguchi, K., see Tanaka, N. 95
Jacobsson, S., see Hagman, A. 117
Jinno, K., see Wahyuni, W. T. 398
Karger, B. L., see Guttman, A. 41
Kaufmann, S. H. E., see Schoel, B. 165
Kiss, L.
——, Tar, A., Gál, S., Toth-Martinez, B. L. and Hernádi, F. J.
Modified general affinity adsorbent for large-scale purification of penicillinases 109
Koike, H., see Arai, T. 439
Korpela, T., see Mäkelä, M. 391
Koyama, Y., see Hashimoto, H. 182
Krupčik, J.
——, Repka, D., Benická, E., Hevesi, T.,. Nolte, J., Paschold, B. and Mayer, H.
Use of retention temperatures for the identification of phenols separated on OV-1701 capillary columns using linear temperature-programmed gas chromatography 203
Lawson, D. E. M., see Okano, T. 145
Lázaro, F.
——, Luque de Castro, M. D. and Valcárcel, M.
Fluorimetric determination of aflatoxins in foodstuffs by high-performance liquid chromatography with flow injection analysis 173
Li, Z., see Grob, K. 387
Lienne, M.
——, Caude, M., Rosset, R. and Tambute, A.
Optimization of the direct chiral separation of potential cytotoxic α-methylene-γ-butyrolactones and α-methylene-γ-butyrolactams by liquid chromatography 55
Luque de Castro, M. D., see Lázaro, F. 173
Mäkelä, M.
——, Mattsson, P., Pintamo-Kenttälä, K. and Korpela, T.
Cyclodextrin–ligand interaction as a simplified model of biospecific affinity chromatography 391
Marston, A., see Slacanin, I. 265
Matsuura, K., see Harada, K.-I. 275
Mattsson, P., see Mäkelä, M. 391
Mayer, H., see Krupčik, J. 203
Montanarella, L., see Gagliardi, L. 296
Morris, M. D., see Peck, K. 193
Nam, H. S.
——, Cho, S. Y. and Rhee, J. S.
High-performance liquid chromatographic analysis of major carotenoids from *Rhodotorula glutinis* 445

Narce, M.
——, Gresti, J. and Bezard, J.
Method for evaluating the bioconversion of radioactive polyunsaturated fatty acids by use of reversed-phase liquid chromatography 249
Nau, H., see Wittfoht, W. 433
Netting, A. G., see Rhodes, S. H. 135
Nicoletti, I., see Polcaro, C. 127
Nolte, J., see Krupčik, J. 203
Nomura, A.
——, Yamada, J. and Tsunoda, K.-I.
Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 87
Novotny, M., see Flurer, C. L. 73
Ohnishi, S., see Tanaka, N. 95
Oishi, S., see Harada, K.-I. 275
Ōizumi, H., see Arai, T. 439
Oka, H., see Harada, K.-I. 275
Okamoto, Y.
——, Aburatani, R. and Hatada, K.
Chromatographic resolution. XXI. Direct optical resolution of abscisic acid by high-performance liquid chromatography on cellulose tris(3,5-dimethylphenylcarbamate) 454
Okano, T.
—— and Lawson, D. E. M.
Chick 28 000 M_r vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 145
Okuno, A., see Tanaka, N. 95
Ossicini, L., see Polcaro, C. 127
Palitti, F., see Polcaro, C. 127
Paschold, B., see Krupčik, J. 203
Pasechnik, V. A., see Gorbunov, A. A. 307
Patrykiejew, A., see Gawdzik, B. 233
Paulus, A., see Guttman, A. 41
Peck, K.
—— and Morris, M. D.
Optical errors in a liquid chromatography absorbance cell 193
Pintamo-Kenttälä, see Mäkelä, M. 391
Polcaro, C.
——, Nicoletti, I., Ossicini, L., Caponecchi, G., Cozzi, R., Fiore, M. and Palitti, F.
Chromatographic and cytogenetic analysis of *in vivo* metabolites of fluoranthene 127
Rebbani, N., see Girard, J. E. 355
Repka, D., see Krupčik, J. 203
Rhee, J. S., see Nam, H. S. 445
Rhodes, S. H.
—— and Netting, A. G.
Normal-phase high-performance liquid chromatography of triacylglycerols 135
Rimmer, S. R., see Dahiya, J. S. 448
Riley, C. M., see Crouch, Jr., F. W. 333
Risch, B.
——, Galensa, R. and Herrmann, K.
Analysis of catechin and epicatechin by high-performance liquid chromatography after benzoylation 291
Rosset, R., see Lienne, M. 55
Rüstig, H., see Voelkel, A. 219
Scaysbrook, T., see Greenaway, W. 284
Schoel, B.
—— and Kaufmann, S. H. E.
Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for β-galactosidase fusion proteins 165
Schwenk, S.
——, Altmayer, B. and Eichhorn, K. W.
Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary gas chromatography 424
Scott, W. J., see Wittfoht, W. 433
Shimamura, T., see Hashimoto, H. 182
Shirai, Y., see Hashimoto, K. 241
Siefermann-Harms, D.
High-performance liquid chromatography of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 411
Slacanin, I.
——, Marston, A. and Hostettmann, K.
High-performance liquid chromatographic determination of molluscicidal saponins from *Phytolacca dodecandra* (phytolaccaceae) 265
Solovyova, L. Ya., see Gorbunov, A. A. 307
Stephens, R. E.
Separation of tubulin subunits by reversed-phase high-performance liquid chromatography 371
Stobaugh, J. F., see Crouch, Jr., F. W. 333
Stromberg, R.
Statistical optimization of a reversed-phase ion-pair liquid chromatographic method for the analysis of tolmetin sodium in dosage forms 1
Suzuki, M., see Harada, K.-I. 275
Szymanowski, J., see Voelkel, A. 219
Takai, N., see Tanaka, N. 95
Tambute, A., see Lienne, M. 55
Tanaka, N.
——, Hashidzume, K., Araki, M., Tsuchiya, H., Okuno, A., Iwaguchi, K., Ohnishi, S. and Takai, N.
Internal structures of wide-pore packing materials for high-performance liquid chromatography studied by transmission electron microscopy 95
Tar, A., see Kiss, L. 109

Tonelli, D., see Gagliardi, L. 296
Toth-Martinez, B. L., see Kiss, L. 109
Tsuchiya, H., see Tanaka, N. 95
Tsunoda, K.-I., see Nomura, A. 87
Valcárcel, M., see Lázaro, F. 173
Vidrich, V., see Heimler, D. 301
Voelkel, A.
——, Szymanowski, J., Beger, J. and Rüstig, H.
Polarity of some individual derivatives of α,ω-diaminooligoethers as measured by gas chromatography 219
Wahyuni, W. T.
—— and Jinno, K.
Separation of tocopherols on various chemically bonded phases in microcolumn liquid chromatography 398
Wang, K.-T., see Chiou, S.-H. 404
Watanabe, M. F., see Harada, K.-I. 275
Whatley, F. R., see Greenaway, W. 284
Wičar, S.
Flow velocity profiles in rectangular channels containing two liquid layers of different densities and viscosities for sedimentation field flow fractionation 456
Wilson, T. D.
Sample solvent effects in an apparent chiral high-performance liquid chromatographic separation on β-cyclodextrin 31
Wittfoht, W.
——, Scott, W. J. and Nau, H.
Assay of methoxyacetic acid in body fluids and tissues by gas chromatography–mass spectrometry following *tert.*-butyldimethylsilylation 433
Wollenweber, E., see Greenaway, W. 284
Yamada, J., see Nomura, A. 87
Yamazaki, M., see Fujii, Y. 157